

## Measure Definitions

The codes provided in the table below are listed as samples that may satisfy the accompanying measure. The list is not comprehensive and does not represent an endorsement of various coding practices.

Measure Name	Measure Definition	MAP	SH	Sample Codes
Adolescent Preventive Care (ADL - QARR)	<p>The percentage of adolescents ages 12-17 who had at least one outpatient visit with a PCP or OB/GYN practitioner during the measurement year and received the following four components of care during the measurement year:</p> <ul style="list-style-type: none"> <li>• Assessment or counseling or education on risk behaviors and preventive actions associated with sexual activity.</li> <li>• Assessment or counseling or education for depression.</li> <li>• Assessment or counseling or education about the risks of tobacco usage.</li> </ul> <p>Assessment or counseling or education about the risks of substance use (including alcohol and excluding tobacco).</p>		X	<p><i>ICD-10 Codes:</i>            Z13.31; Z71.7;            Z70.3; Z70.9;            Z30.0; Z30.01;            Z71.6; Z71.41;            Z71.51</p>
Adult BMI Assessment (ABA)	The percentage of members 18–74 years of age who had an outpatient visit and whose body mass index (BMI) was documented during the measurement year or the year prior to the measurement year.	X	X	<p><i>ICD-10 Codes:</i>            Z68.41; Z68.43;            Z68.45</p>
Antidepressant Medication Management (AMM)	Number of adults 18 years of age and older with a diagnosis of major depression who were newly treated with antidepressant medication and remained on their antidepressant medications.		X	Pharmaceutical Prescription – see Appendix 1 for sample medications
Asthma Medication Ratio (AMR)	Number of adults and children 5–64 years of age who were identified as having persistent asthma and had a ratio of controller medications to total asthma medications of 0.50 or greater during the measurement year.		X	Pharmaceutical Prescription – see Appendix 1 for sample medications
Breast Cancer Screening (BCS)	Percentage of women 50–74 years of age who had at least one	X	X	<p><i>CPT Codes:</i>            77055-77057</p>

	mammogram to screen for breast cancer in the past two years.			77061-77063 <i>Exclusion Codes:</i> 19180; 19200
Care for Older Adults – Advance Care Planning (COA)	The percentage of adults 66 years and older who had advance care planning during the measurement year.	X		<i>CPT Codes:</i> 99483; 99497
Care for Older Adults – Functional Status Assessment (COA)	The percentage of adults 66 years and older who had a functional status assessment during the measurement year.	X		<i>CPT II Code:</i> 1170F
Care for Older Adults – Medication Review (COA)	The percentage of adults 66 years and older who had a medication review during the measurement year.	X		<i>CPT Codes:</i> 90863; 99483; 99605; 99606
Care for Older Adults – Pain Assessment (COA)	The percentage of adults 66 years and older who had a pain assessment during the measurement year.	X		<i>CPT II Codes:</i> 1125F; 1126F
Cervical Cancer Screening (CCS)	Percentage of women 21–64 years of age who were screened for cervical cancer using either of the following criteria: <ul style="list-style-type: none"> <li>• Women age 21–64 who had cervical cytology performed every 3 years.</li> <li>• Women age 30–64 who had cervical cytology/human papillomavirus (HPV) co-testing performed every 5 years.</li> </ul>		X	<i>CPT Codes:</i> 88141-88143 88152-88154 87620-87622; 51925; 56308; 57540
Colorectal Screening (COL)	Percentage of adults 50–75 who had appropriate screening for colorectal cancer with any of the following tests: <ul style="list-style-type: none"> <li>• 1 Year: Fecal Occult Blood Test.</li> <li>• 3 Years: Stool DNA Test.</li> <li>• 5 Years: Flexible Sigmoidoscopy.</li> <li>• 5 Years: Computed Tomography Colonography.</li> <li>• 10 Years: Colonoscopy.</li> </ul>	X	X	<i>CPT Codes:</i> 44388-44394; 74261-74263; 81528; 82270; 82274; 45330- 45335 <i>Exclusion Codes:</i> 44150-44153 44155-44158
Comprehensive Diabetes Care Eye Exam (CDC)	Percentage of adults 18–75 years of age with diabetes (type 1 and type 2) who had an eye exam (retinal) performed.	X	X	<i>CPT II Codes:</i> 3072F; 2022F; 2023F
Comprehensive Diabetes Care HbA1c Results ((CDC) including control and poor control)	All HbA1c test results in order to evaluate the percentage of adults 18–75 years of age with diabetes (type 1 and type 2) who had HbA1c control (<8.0%) and poor control (>9.0%). A lower score is preferred.	X	X	<i>CPT II Codes:</i> 3045F; 3046F; 3051F; 3052F

Comprehensive Diabetes Care HbA1c Test (CDC)	Percentage of adults 18–75 years of age with diabetes (type 1 and type 2) who had hemoglobin A1c (HbA1c) testing.		X	<i>CPT Codes:</i> 83036
Comprehensive Diabetes Care Nephropathy Screening (CDC)	Percentage of adults 18–75 years of age with diabetes (type 1 and type 2) who had Medical attention for nephropathy.	X	X	<i>CPT II Codes:</i> 3066F 4010F  <i>Exclusion Codes:</i> 50380; 50340; 50370
Controlling High Blood Pressure < 140/90 (CBP)	Percentage of adults 18–85 years of age who had a diagnosis of hypertension and whose blood pressure was adequately controlled.	X	X	<i>CPT II Codes:</i> 3080F; 3078F
Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medication (SSD)	Assesses adults 18–64 years of age with schizophrenia or bipolar disorder, who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.		X	Pharmaceutical Prescription – see Appendix 1 for sample medications
HIV Viral Load Suppression (VLS – QARR)	The percentage of Medicaid enrollees confirmed HIV-positive who had a HIV viral load less than 200 copies/mL at last HIV viral load test during the measurement year.		X	N/A
Medication Reconciliation Post-Discharge (MRP)	The percentage of discharges from January 1–December 1 of the measurement year for members 18 years of age and older for whom medications were reconciled the date of discharge through 30 days after discharge (31 total days).	X		<i>CPT Codes:</i> 99483; 99495; 99496  <i>CPT II Codes:</i> 1111F
Sexually Transmitted Infections: Screening for Chlamydia (STI)	Patients with at least one urine/cervical/urethral test for Chlamydia during measurement year.		X	<i>CPT Codes:</i> 86631; 86632 <i>HCPCS Codes:</i> G9228; G9229
Sexually Transmitted Infections: Screening for Gonorrhea (STI)	Patients with at least one urine/cervical/urethral test for Gonorrhea during the measurement year.		X	<i>CPT Codes:</i> 87590; 87591 <i>HCPCS Codes:</i> G9228; G9229
Sexually Transmitted Infections: Screening for Syphilis (STI)	Patients with Syphilis screening performed during measurement year.		X	<i>CPT Codes:</i> 86592; 86593 <i>HCPCS Codes:</i> G9228; G9229

<p>Use of Spirometry Testing in the Assessment and Diagnosis of COPD (SPR)</p>	<p>Assesses adults 40 years of age and older who have a new diagnosis of chronic obstructive pulmonary disease (COPD) or newly active COPD, who received spirometry testing to confirm the diagnosis.</p>		<p>X</p>	<p><i>CPT Codes:</i> 94010; 94014; 94015; 94016; 94060</p>
<p>Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC)</p>	<p>Assesses children and adolescents 3-17 years of age who had an outpatient visit with a primary care practitioner or OB/GYN during the measurement year and had evidence of:</p> <ul style="list-style-type: none"> <li>• Body mass index (BMI) percentile documentation.</li> <li>• Counseling for nutrition.</li> <li>• Counseling for physical activity.</li> </ul>		<p>X</p>	<p><i>BMI Codes:</i> N/A</p> <p><i>Nutrition CPT Codes:</i> 97802; 97803</p> <p><i>Physical Activity ICD-10 CM Codes:</i> Z02.5; Z71.82</p>