

# HEDIS® & QARR Provider Reference Guide- MY 2022



## Table of Contents

<b>Glossary of Terms:</b> .....	5
<b>Effectiveness of Care- <i>Prevention and Screening Measures:</i></b> .....	6
Weight Assessment & Counseling for Nutrition and Physical Activity for Children & Adolescents .....	6
Childhood Immunizations Status .....	6
Immunizations for Adolescents .....	6
Lead Screening in Children.....	6
Breast Cancer Screening .....	6
Cervical Cancer Screening.....	7
Colorectal Cancer Screening .....	7
Chlamydia Screening in Women .....	7
Care of Older Adults.....	7
Flu Shots for Adults Ages 18-64 .....	7
Flu Shots for Adults Ages 65 and Older .....	8
Pneumococcal Vaccination Status for Older Adults .....	8
Medical Assistance with Smoking and Tobacco Cessation .....	8
<b>Effectiveness of Care- <i>Respiratory Conditions Measures:</i></b> .....	9
Appropriate Testing for Pharyngitis.....	9
Use of Spirometry Testing in the Assessment and Diagnosis of COPD.....	9
Pharmacotherapy Management of COPD Exacerbation .....	9
Asthma Medication Ratio .....	9
<b>Effectiveness of Care- <i>Cardiovascular Conditions Measures:</i></b> .....	11
Controlling High Blood Pressure .....	11

Persistence of Beta-Blocker Treatment After a Heart Attack..... 11

Statin Therapy for Patients with Cardiovascular Disease ..... 11

Cardiac Rehabilitation ..... 12

**Effectiveness of Care- *Diabetes Measures:*** ..... 13

    Hemoglobin A1c Control for Patients with Diabetes ..... 13

    Blood Pressure Control for Patients with Diabetes ..... 13

    Eye Exam for Patients with Diabetes ..... 13

    Kidney Health Evaluation for Patients with Diabetes ..... 14

    Statin Therapy for Patients with Diabetes ..... 14

**Effectiveness of Care- *Musculoskeletal Conditions Measures:***..... 16

    Osteoporosis Management in Women Who Had a Fracture ..... 16

    Osteoporosis Screening in Older Women..... 16

**Effectiveness of Care- *Behavioral Health Measures:*** ..... 17

    Diagnoses of Mental Health Disorders ..... 17

    Antidepressant Medication Management..... 17

    Follow-up Care for Children Prescribed ADHD Medication ..... 17

    Follow-Up After Hospitalization for Mental Illness..... 17

    Follow-Up After Emergency Department Visit for Mental Illness ..... 18

    Diagnosed Substance Use Disorders..... 18

    Follow-Up After High-Intensity Care for Substance Use Disorder ..... 18

    Follow-Up After Emergency Department Visit for Substance Use ..... 18

    Pharmacotherapy for Opioid Use Disorder ..... 19

    Initiation of Pharmacotherapy Upon New Episode of Opioid Dependence ..... 19

    Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications..... 19

Diabetes Monitoring for People with Diabetes and Schizophrenia..... 19

Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia ..... 20

Adherence to Antipsychotic Medications for Individuals with Schizophrenia ..... 20

Metabolic Monitoring for Children and Adolescents on Antipsychotics..... 20

**Effectiveness of Care- Care Coordination Measures:**..... 21

    Advance Care Planning ..... 21

    Transitions of Care ..... 21

    Follow-Up After Emergency Department Visit for People with Multiple High-Risk Chronic Conditions..... 21

**HIV Care** ..... 22

    Viral Load Suppression..... 22

    Sexually Transmitted Diseases: Chlamydia Screening ..... 22

    Sexually Transmitted Diseases: Gonorrhea Screening ..... 22

    Sexually Transmitted Diseases: Syphilis Screening ..... 22

**Access/Availability of Care Measures:** ..... 23

    Adults’ Access to Preventative/Ambulatory Health Services ..... 23

    Annual Dental Visit..... 23

    Initiation and Engagement of Substance Use Disorder Treatment ..... 23

    Prenatal and Postpartum Care..... 23

    Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics..... 24

**Utilization and Risk Adjusted Utilization Measures:**..... 25

    Well-Child Visits in the First 30 Months of Life..... 25

    Child and Adolescent Well-Care Visits ..... 25

**References and Comments:** ..... 26

## Glossary of Terms:

Measure Name: The name of the HEDIS measure used for reporting

Measure Code: The code used to identify the measure name

Age Criteria: The age of the population that is eligible for the measure

Measure Description: The description of what is being measured in each domain

ICD10 Codes: The diagnosis code assigned to conditions and diseases used for claims reporting

CPT Codes: Codes used to describe the treatment being provided

HCPCS Codes: Codes used to describe supplies, equipment and devices being provided

LOINC Codes: Codes use to describe the tests being provided, such as laboratory tests and vital signs

Exclusions: Some measures remove eligible members from a measure they are otherwise eligible for (ex: hospice, palliative care, ESRD/dialysis)

Measurement Year (MY): Refers to the current calendar year (January 1, 2022 – December 31, 2022), for the measures unless otherwise specified

### Effectiveness of Care- *Prevention and Screening Measures:*

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Weight Assessment & Counseling for Nutrition and Physical Activity for Children & Adolescents ( <b>WCC</b> )	3-17 years of age	Eligible members with an outpatient PCP and/or OB/GYN visit with documentation of BMI profile and counseling for nutrition and physical activity, within the MY	<b>ICD 10 Codes:</b> Z68.51-Z68.54, Z02.5 <b>CPT Codes:</b> 97802-97804, 99201-99205 <b>Exclusion:</b> Members on/using hospice services (99377); Exclusion for BMI only- females with a pregnancy diagnosis during the MY	Select Health
Childhood Immunizations Status ( <b>CIS</b> )	2 years of age	Children who have received four doses of <b>DTaP</b> (diphtheria, tetanus, and pertussis) and <b>PCV</b> (pneumococcal conjugate); three doses of <b>IPV</b> (polio), <b>HepB</b> (hepatitis B), and <b>HIB</b> (haemophilus influenza B), two to three doses of <b>RV</b> (rotavirus), two Flu vaccines (influenza), and one dose of <b>MMR</b> (measles, mumps, and rubella), <b>VZV</b> (chicken pox) and <b>HepA</b> (Hepatitis A) by their second birthday	<b>ICD 10 Codes:</b> B15.0, B16.0, B01.0, B26.0 <b>CPT Codes:</b> 90697, 90647, 90633, 90707, 90670, 90710 <b>Exclusion:</b> Members on/using hospice services (99377), diagnosis of HIV (B20, B97.35) and immunocompromised conditions (D80.0)	Select Health
Immunizations for Adolescents ( <b>IMA</b> )	13 years of age	Adolescents who have received one dose of Tdap (tetanus, diphtheria, and pertussis) and meningococcal vaccines, and have had the HPV (human papillomavirus) vaccine series finalized, by their 13 <sup>th</sup> birthday.	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 90649, 90619, 90715 <b>Exclusions:</b> Members on/using hospice services (99377)	Select Health
Lead Screening in Children ( <b>LSC</b> )	2 years of age	Children who have completed at least one lead blood test (capillary or venous) by their second birthday, to determine lead poisoning	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 8365 <b>LOINC Codes:</b> 5671-3 <b>Exclusions:</b> Members on/using hospice services (99377)	Select Health
Breast Cancer Screening ( <b>BCS</b> )	Women 52-74 years of age	Women between 50 to 74 years of age, who have had a mammogram within 2 years of the MY, to screen for breast cancer	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 99221, 99291, 77061, 99211 <b>Exclusions:</b> Members on/using hospice services (9937) or palliative care (Z51.5), and those who have either a unilateral (19180) or bilateral mastectomy (Z90.13) modifier (optional), members 66 and older residing in a facility or members 66 and older with a	Select Health, MAP Total, EasyCare and EasyCare Plus

HEDIS® & Quality Assurance Reporting Requirements (QARR) Provider Reference Guide- MY2022

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
			diagnosis frailty (L89.000) <b>and</b> advanced illness (A81.00)	
Cervical Cancer Screening (CCS)	Women 24-64 years of age	Women 21 to 64 years of age, who have had a cervical cytology (21 to 64) within 3 years, cervical high-risk human papillomavirus (30-64) within 5 years or both contesting (30-64) within the last 5 years, to screen for cervical cancer	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 88141-88143, 87624, 87625 <b>Exclusions:</b> Members on/using hospice services (G9473) or palliative care (Z51.5) or have had a hysterectomy (51925) and/or absence of cervix (Q51.5) (optional)	Select Health
Colorectal Cancer Screening (COL)	51-75 years of age	Members 50 to 75 years of age, who have had at least one of the following cancer screenings within the MY and prior year: Fecal Occult Blood Test (FOBT), Flexible Sigmoidoscopy, Colonoscopy, CT Colonography and/or FIT-DNA testing	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 82270, 45330, 44388, 74261, 81528 <b>Exclusions:</b> Members on/using hospice services (9937) or palliative care (Z51.5), members 66 and older residing in a facility or members 66 and older with a diagnosis frailty (L89.000) <b>and</b> advanced illness (A81.00), and members with a diagnosis of colorectal cancer (C18.0) or total colectomy (44150) (optional)	MAP Total, EasyCare and EasyCare Plus
Chlamydia Screening in Women (CHL)	Women 16-24 years of age	Sexually active women between 16 to 24 years of age, who have at least one test for chlamydia within the MY	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 87110, 87320, 70010, 70015 <b>Exclusions:</b> Members on/using hospice services (9937), and members who have had a pregnancy test (81025) with a retinoid prescription <b>or</b> radiology diagnostic test (70010) at time of test	Select Health
Care of Older Adults (COA)	66 years of age	Members who are 66 years of age who have completed all three at least once within the MY- Pain and functional status assessments, as well as medication review (including OTC and supplements)	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 99843, 90863, 99495 <b>HCPCS Codes:</b> G8427, G0438 <b>Exclusions:</b> Members on/using hospice services (9937), services provided in an acute inpatient setting (99221)	MAP Total, EasyCare and EasyCare Plus
Flu Shots for Adults Ages 18-64 (FVA) *CAHPS	18-64 years of age	Members between 18 to 64 years of age, who have received a flu vaccine or flu spray, between July 1 <sup>st</sup> and December 31 <sup>st</sup> of the MY	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> NA <b>Exclusions:</b> NA	Select Health

HEDIS® & Quality Assurance Reporting Requirements (QARR) Provider Reference Guide- MY2022

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Flu Shots for Adults Ages 65 and Older <b>(FVO)</b> <b>*CAHPS</b>	65 years of age and older	Members 65 years of age and older, who have received a flu vaccine between July 1 <sup>st</sup> and December 31 <sup>st</sup> of the MY	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> NA <b>Exclusions:</b> NA	MAP Total, EasyCare and EasyCare Plus
Pneumococcal Vaccination Status for Older Adults <b>(PNU)</b> <b>*CAHPS</b>	65 years of age and older	Members 65 years of age and older, who have received at least one pneumococcal vaccine	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> NA <b>Exclusions:</b> NA	MAP Total, EasyCare and EasyCare Plus
Medical Assistance with Smoking and Tobacco Cessation <b>(MSC)</b> <b>*CAHPS</b>	18 years of age and older	Members 18 years of age or older, who have medical assistance with advising smokers/tobacco users to quit, as well as cessation medications and strategies during the MY	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> NA <b>Exclusions:</b> NA	Select Health, MAP Total, EasyCare and EasyCare Plus



### Effectiveness of Care- Respiratory Conditions Measures:

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Appropriate Testing for Pharyngitis (CWP)	3 years of age and older	Members 3 years of age and older, who were tested and diagnosed with pharyngitis, given an antibiotic and received a strep test (group A streptococcus) <i>Age groups include: 3-17, 18-64 and 65 years and older</i>	<b>ICD 10 Codes:</b> J02.0, J02.8, J03.01 <b>CPT Codes:</b> 87070, 87081, 87430 <b>Exclusions:</b> Members on/using hospice services (9937), comorbid conditions within the 12 months prior to exam (44.0, Z21), and members on antibiotics within 30 days prior to exam.	Select Health, MAP Total, EasyCare and EasyCare Plus
Use of Spirometry Testing in the Assessment and Diagnosis of COPD (SPR)	42 years of age or older	Members 40 years of age or older, who received adequate spirometry testing to determine a new diagnosis of COPD, within the MY and the year prior	<b>ICD 10 Codes:</b> J44.0, J44.1, J.44.9 <b>CPT Codes:</b> 94010, 94014, 94060, 94375 <b>Exclusions:</b> Members on/using hospice services (9937), members with a previously determined diagnosis of COPD (J44.0), emphysema (J43.0) or chronic bronchitis (J41.0)	Select Health, MAP Total, EasyCare and EasyCare Plus
Pharmacotherapy Management of COPD Exacerbation (PCE)	40 years of age	Members 40 years of age, that were hospitalized or had an emergency room visit between January 1 <sup>st</sup> and November 30 <sup>th</sup> , with COPD exacerbation, who were prescribed appropriate medications (a systemic corticosteroid or a bronchodilator) <i>ED visits resulting in a hospitalization are not included</i>	<b>ICD 10 Codes:</b> J44.0, J44.9, J43.0, J43.9 <b>CPT Codes:</b> 99281-99285 <b>Exclusions:</b> Members on/using hospice services (9937), and nonacute hospitalizations	Select Health, MAP Total, EasyCare and EasyCare Plus
Asthma Medication Ratio (AMR)	5-64 years of age	Members 5 to 64 years of age, with a diagnosis of persistent asthma that have controller medications vs total asthma medications prescribed ratio of at least 0.50 <i>Age groups include: 5-11, 12-18, 19-50, 51-64 years</i>	<b>ICD 10 Codes:</b> J45.21, J45.51, J45.991 <b>CPT Codes:</b> 99221-99223, 99281, 98969, 99201-99205, 98966 <b>Exclusions:</b> Members on/using hospice services (9937), members who had a diagnosis of emphysema (J43.0), COPD	Select Health

HEDIS® & Quality Assurance Reporting Requirements (QARR) Provider Reference Guide- MY2022

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
			(J44.0), obstructive chronic bronchitis, chronic respiratory conditions (J68.4), cystic fibrosis and acute respiratory failure (J96.00), and members without any asthma controlling medications	

### Effectiveness of Care- *Cardiovascular Conditions Measures:*

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Controlling High Blood Pressure (CBP)	18-85 years of age	Members 18 to 85 years of age with a diagnosis of HTN (hypertension) whose most recent blood pressure measurements, during the MY, show an adequate control of condition, as evidenced by results of <140/90mm Hg	<p><b>ICD 10 Codes:</b> I10</p> <p><b>CPT Codes:</b> NA</p> <p><b>LOIN Codes:</b> Diastolic- 8462-4; Systolic- 8480-6</p> <p><b>Exclusions:</b> Members on/using hospice services (9937) or palliative care (Z51.5), members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) <b>and</b> advanced illness (A81.00) diagnosis, and members 81 and older with a frailty (L89.000) diagnosis. Optional exclusions include members with a diagnosis of End-stage renal disease (ESRD) (N18.5), nephrectomy (50220) or kidney transplant (50360), and female members who are pregnant (O00.0)</p>	Select Health, MAP Total, EasyCare and EasyCare Plus
Persistence of Beta-Blocker Treatment After a Heart Attack (PBH)	18 years of age and older	Members 18 years of age and older who were discharged from a hospitalization for AMI (acute myocardial infarction), that were prescribed a six-month treatment of persistent beta-blockers	<p><b>ICD 10 Codes:</b> I21.01, I21.19, I21.29</p> <p><b>CPT Codes:</b> 99221-99223, 99232, 99253</p> <p><b>Exclusions:</b> Members on/using hospice services (9937), members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) <b>and</b> advanced illness (A81.00) diagnosis, and members 81 and older with a frailty (L89.000) diagnosis. Optional exclusions include members diagnosed with asthma (J45.21), COPD (J44.0), obstructive chronic bronchitis, chronic respiratory conditions (J68.4) and hypotension.</p>	Select Health, MAP Total, EasyCare and EasyCare Plus
Statin Therapy for Patients with Cardiovascular Disease (SPC)	Males 21-75 years of age; Females 40-	Male members between 21 to 75 years of age, and female members between 40 to 75 years of age, who have had an ASCVD diagnosis (atherosclerotic cardiovascular	<p><b>ICD 10 Codes:</b> I21.01, I21.9, I25.2</p> <p><b>CPT Codes:</b> 99221-99223, 33510, 33513</p> <p><b>Exclusions:</b> Members on/using hospice services (9937) or palliative care (Z51.5),</p>	Select Health, MAP Total,

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
	75 years of age	disease) and has received statin therapy, as well as maintained an 80% of statin adherence	members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) <b>and</b> advanced illness (A81.00) diagnosis, members diagnosed with ESRD (N18.5), cirrhosis (K70.30), myalgia (M79.1), myositis (M60.80), myopathy (G72.9) or rhabdomyolysis (M62.82) and female members with a pregnancy (O00.0) diagnosis, or received IVF (in vitro fertilization) (HCPCS S4015) or prescribed clomiphene during the MY	EasyCare and EasyCare Plus
Cardiac Rehabilitation (CRE)	18 years of age and older	Members 18 years of age and older, who participated in cardiac rehab post qualifying cardiac event (such as an MI/CABG). <i>Rates reported include initiation (2 or more sessions within 30 days), engagement 1 (12 or more sessions within 90 days), engagement 2 (24 or more sessions within 180 days) and achievement (36 or more sessions within 180 days)</i>	<b>ICD 10 Codes:</b> I21.01, I21.9, I25.2 <b>CPT Codes:</b> 93797, 93798 <b>Exclusions:</b> Members on/using hospice services (9937) or palliative care (Z51.5), members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) <b>and</b> advanced illness (A81.00) diagnosis, members 81 and older with a frailty (L89.000) diagnosis. Members diagnosed with an MI (I21.01), CABG (33510), heart or heart & lung transplant (33927) or heart valve repair/replacement (33361) within 180 days after initial event are also excluded.	Select Health, MAP Total, EasyCare and EasyCare Plus

### Effectiveness of Care- *Diabetes Measures:*

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Hemoglobin A1c Control for Patients with Diabetes (HBD)	18-75 years of age	Members between 18 to 75 years of age, diagnosed with diabetes (types 1 and 2), who have a completed hemoglobin A1c within the MY.  <i>A1c levels &lt;8.0 indicate good control and levels &gt;9.0 indicate poor control.</i>	<b>ICD 10 Codes:</b> E10.10, E10.36, E11.00 <b>CPT Codes:</b> 83036, 83067 <b>Exclusions:</b> Members on/using hospice services (9937) or palliative care (Z51.5), members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) <b>and</b> advanced illness (A81.00) diagnosis, members without a diagnosis of diabetes, members diagnosed with polycystic ovarian syndrome (E28.2), gestational (O24.419) or steroid (E09.9) induced diabetes	Select Health, MAP Total, EasyCare and EasyCare Plus
Blood Pressure Control for Patients with Diabetes (BPD)	18-75 years of age	Members between 18 to 75 years of age, diagnosed with diabetes (types 1 and 2), whose most recent blood pressure measurements, during the MY, show an adequate control of condition, as evidenced by results of <140/90mm Hg	<b>ICD 10 Codes:</b> E10.10, E10.36, E11.00 <b>CPT Codes:</b> 99221, 99281, 99217,99201 <b>LOINC Codes:</b> Diastolic- 9462-4, Systolic- 8480-6 <b>Exclusions:</b> Members on/using hospice services (9937) or palliative care (Z51.5), members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) <b>and</b> advanced illness (A81.00) diagnosis, members without a diagnosis of diabetes, members diagnosed with polycystic ovarian syndrome (E28.2), gestational (O24.419) or steroid (E09.9) induced diabetes	Select Health, MAP Total, EasyCare and EasyCare Plus
Eye Exam for Patients with Diabetes (EED)	18-75 years of age	Members between 18 to 75 years of age, diagnosed with diabetes (types 1 and 2), who have completed a retinal eye exam	<b>ICD 10 Codes:</b> E10.10, E10.36, E11.00 <b>CPT Codes:</b> 60278, 67041, 67210, 92229 <b>Exclusions:</b> Members on/using hospice services (9937) or palliative care (Z51.5),	Select Health, MAP Total,

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
			members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) <b>and</b> advanced illness (A81.00) diagnosis, members without a diagnosis of diabetes and diagnosed with polycystic ovarian syndrome (E28.2), gestational (O24.419) or steroid (E09.9) induced diabetes	EasyCare and EasyCare Plus
Kidney Health Evaluation for Patients with Diabetes (KED)	18-85 years of age	Members between 18 to 85 years of age, diagnosed with diabetes (types 1 and 2), who completed the eGFR (estimated glomerular filtration rate) <b>and</b> uACR (urine albumin-creatinine ratio) lab tests during the MY. <i>*The uACR must have both a quantitative urine albumin test <b>and</b> a urine creatinine test to be complete</i>	<b>ICD 10 Codes:</b> E10.10, E10.36, E11.00 <b>CPT Codes:</b> 99221, 82043, 82570, 80047 <b>LOINC Codes:</b> 13705-9, 48642-3 <b>Exclusions:</b> Members on/using hospice services (9937) or palliative care (Z51.5), members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) <b>and</b> advanced illness (A81.00) diagnosis, members 81 and older with a frailty (L89.000) diagnosis, members diagnosed with ESRD (N18.5) or on dialysis, and members. Optional exclusions include members without a diagnosis of diabetes and diagnosed with polycystic ovarian syndrome (E28.2), gestational (O24.419) or steroid (E09.9) induced diabetes	Select Health, MAP Total, EasyCare and EasyCare Plus
Statin Therapy for Patients with Diabetes (SPD)	40-75 years of age	Members between 40 to 75 years of age, with a diagnosis of diabetes, who do not have a ASCVD diagnosis (atherosclerotic cardiovascular disease), and has received statin therapy, as well as maintained an 80% of statin adherence	<b>ICD 10 Codes:</b> E10.10, E10.36, E11.00 <b>CPT Codes:</b> 99221, 99281, 99217, 99201 <b>Exclusions:</b> Members on/using hospice services (9937) or palliative care (Z51.5), members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) <b>and</b> advanced illness (A81.00) diagnosis, members diagnosed with a	Select Health, MAP Total, EasyCare and EasyCare Plus

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
			cardiovascular or ischemic vascular disease, ESRD (N18.5), cirrhosis (K70.30), myalgia (M79.1), myositis (M60.80), myopathy (G72.9) or rhabdomyolysis (M62.82) and female members with a pregnancy (O00.0) diagnosis, or received IVF (in vitro fertilization) (HCPCS S4015) or prescribed clomiphene during the MY. Optional exclusions include members without a diagnosis of diabetes and diagnosed with polycystic ovarian syndrome (E28.2), gestational (O24.419) or steroid (E09.9) induced diabetes	

### Effectiveness of Care- Musculoskeletal Conditions Measures:

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Osteoporosis Management in Women Who Had a Fracture <b>(OMW)</b>	67-85 years of age	Female members between 67 to 85 years of age, who have had a BMD (bone mineral density) test or prescribed a medication for osteoporosis within six months, after suffering a fracture	<b>ICD 10 Codes:</b> M80.00XA, M80.80XA <b>CPT Codes:</b> 21811, 21820, 23620 <b>HCPCS Codes:</b> J0897, J1740, J3110, J3489 <b>Exclusions:</b> Members on/using hospice services (9937) or palliative care (Z51.5), members 67 years of age or older residing in a facility, members 67 to 80 years of age with a frailty (L89.000) <b>and</b> advanced illness (A81.00) diagnosis, members 81 and older with a frailty (L89.000) diagnosis, and members with a BMD test within 2 years prior to fracture event, had claims for osteoporosis medications within 1 year prior to fracture event. Fractures of the finger, toe, face, and skull are also excluded.	MAP Total, EasyCare and EasyCare Plus
Osteoporosis Screening in Older Women <b>(OSW)</b>	66-75 years of age	Female members between 65-75 years of age, who have been assessed for osteoporosis during the MY and the year prior	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 76977, 77078, 77081, 77085 <b>Exclusions:</b> Members on/using hospice services (9937) or palliative care (Z51.5), members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) <b>and</b> advanced illness (A81.00) diagnosis, and members who have claims for osteoporosis medications prior to the MY or received osteoporosis medications within the 3 years prior to MY	MAP Total, EasyCare and EasyCare Plus



### Effectiveness of Care- Behavioral Health Measures:

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Diagnoses of Mental Health Disorders (DMH)	1 years of age and older	Members 1 years of age and older, who have been diagnosed with a mental health disorder during the MY <i>Age stratifications are 1-17, 18-64 and 65 years of age and older</i>	<b>ICD 10 Codes:</b> F03.90, F21, F31.0, F43.10 <b>CPT Codes:</b> NA <b>Exclusions:</b> Members on/using hospice services (9937)	Select Health, MAP Total, EasyCare and EasyCare Plus
Antidepressant Medication Management (AMM)	18 years of age and older	Members 18 years of age and older with a diagnosis of major depression, who received and remained on antidepressant medications <i>Rates reported include the effective acute phase- those who received medication for at least 84 days, and the effective continuation phase- those who received medication for at least 180 days</i>	<b>ICD 10 Codes:</b> F32.0-F32.4, F32.9, F33.0-F33.3 <b>CPT Codes:</b> 99221, 98960, 90791, 98966 <b>Exclusions:</b> Members on/using hospice services (9937) and members without a diagnosis of major depression	Select Health, MAP Total, EasyCare and EasyCare Plus
Follow-up Care for Children Prescribed ADHD Medication (ADD)	6-12 years of age	Members between 6 to 12 years of age, who were prescribed ADHD (attention deficit/hyperactivity disorder) medications and had a follow-up visit within 30 days of prescription ( <i>initiation phase</i> ) and at least two more follow-up visits within 10 months of prescription ( <i>continuation and maintenance phase</i> )	<b>ICD 10 Codes:</b> F90.0-F90.2, F90.8, F90.9 <b>CPT Codes:</b> 96150, 98960, 99217, 90791 <b>Exclusions:</b> Members on/using hospice services (9937) and members with a diagnosis of narcolepsy (G47.411)	Select Health
Follow-Up After Hospitalization for Mental Illness (FUH)	6 years of age and older	Members 6 years of age and older, who had a mental illness or self-harm diagnosed hospital admission, that was discharged and had a mental health provider follow-up. Follow-up appointments are rated by those who complete an appointment within 7 days	<b>ICD 10 Codes:</b> T14.91XA, F20.0, F31.0, F32.9 <b>CPT Codes:</b> 98960, 98966, 99495, 90791 <b>Exclusions:</b> Members on/using hospice services (9937), any members discharged and readmitted within 30 days	Select Health, MAP Total, EasyCare and

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
		post-discharge, and those who complete an appointment within 30 days post-discharge		EasyCare Plus
Follow-Up After Emergency Department Visit for Mental Illness (FUM)	6 years of age and older	Members 6 years of age and older, who had a mental illness or self-harm diagnosed emergency department visit, that was discharged and had a mental health provider follow-up. Follow-up appointments are rated by those who complete an appointment within 7 days post-discharge, and those who complete an appointment within 30 days post-discharge	<b>ICD 10 Codes:</b> T14.91XA, F20.0, F31.0, F32.9 <b>CPT Codes:</b> 98960, 98966, 99495, 90791 <b>Exclusions:</b> Members on/using hospice services (9937), and members seen in the emergency department resulting in a hospital admission or an admission within 30 days of initial emergency department visit regardless of admission diagnosis	Select Health, MAP Total, EasyCare and EasyCare Plus
Diagnosed Substance Use Disorders (DSU)	13 years of age and older	Members 13 years of age and older with a substance use disorder during the MY. Substance abuse disorders include alcohol, opioid, substance use and other/unspecified drugs	<b>ICD 10 Codes:</b> F10.10, F11.10, F14.10, F15.10 <b>CPT Codes:</b> NA <b>Exclusions:</b> Members on/using hospice services (9937)	Select Health, MAP Total, EasyCare and EasyCare Plus
Follow-Up After High-Intensity Care for Substance Use Disorder (FUI)	13 years of age and older	Members 13 years of age and older with a substance use disorder, that had an acute inpatient hospitalization, residential treatment, or detoxification visit, resulting in follow-up visits. Follow-up appointments are rated by those who complete an appointment within 7 days post-discharge, and those who complete an appointment within 30 days post-discharge	<b>ICD 10 Codes:</b> F10.10, F11.10, F14.10, F15.10 <b>CPT Codes:</b> 99408, 99409 <b>HCPCS Codes:</b> H0017-H0019, H0008-H0009 <b>Exclusions:</b> Members on/using hospice services (9937), any nonacute inpatient admissions other than behavioral health	Select Health, MAP Total, EasyCare and EasyCare Plus
Follow-Up After Emergency	13 years of age and older	Members 13 years of age and older that had an emergency department visit with a primary substance use disorder or drug	<b>ICD 10 Codes:</b> T40.0X1A, F10.10, F11.10, F14.10, F15.10 <b>CPT Codes:</b> 99408, 99409	Select Health,

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Department Visit for Substance Use (FUA)		overdose diagnosis and completed a follow-up visit. Follow-up appointments are rated by those who complete an appointment within 7 days post-discharge, and those who complete an appointment within 30 days post-discharge	<b>HCPCS Codes:</b> G0177, H0020 <b>Exclusions:</b> Members on/using hospice services (9937), and members seen in the emergency department resulting in a hospital admission or an admission within 30 days of initial emergency department visit regardless of admission diagnosis	MAP Total, EasyCare and EasyCare Plus
Pharmacotherapy for Opioid Use Disorder (POD)	16 years of age and older	Members 16 years of age and older with a new opioid use disorder, who receives pharmacotherapy for at least 180 days	<b>ICD 10 Codes:</b> F11.10, F11.20, <b>CPT Codes:</b> NA <b>HCPCS Codes:</b> G2070, G2069, J0572, H0033, H0020, J2315 <b>Exclusions:</b> Members on/using hospice services (9937), and members who had an inpatient stay for 8 or more days during treatment period	Select Health, MAP Total, EasyCare and EasyCare Plus
Initiation of Pharmacotherapy Upon New Episode of Opioid Dependence (POD-N)	18 years of age and older	Members 18 years of age and older, who begin pharmacotherapy for a new opioid dependence treatment within 30 days of new diagnosis	ICD 10 Codes: CPT Codes: Exclusions: Members who are not diagnosed with a new opioid dependence	Select Health
Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications (SSD)	16-64 years of age	Members between 16 and 64 years of age, diagnosed with schizophrenia, or schizoaffective or bipolar disorder with a completed diabetes screening, that were prescribed antipsychotic medications during the MY	<b>ICD 10 Codes:</b> F31.9, F20.9, F25.9 <b>CPT Codes:</b> 80047, 80053, 83036, 83037 <b>Exclusions:</b> Members on/using hospice services (9937), and members who are diagnosed with diabetes or received insulin or oral hypoglycemic/antihyperglycemic medications during the MY or the prior year	Select Health
Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD)	18-64 years of age	Members between 18 and 64 years of age, with a diagnosis of diabetes and schizophrenia or schizoaffective disorder, who completed an HbA1c and LDL-C test during the MY	<b>ICD 10 Codes:</b> E10.9, E11.9, F20.9, F25.9 <b>CPT Codes:</b> 83036, 83037, 80061, 83700 <b>Exclusions:</b> Members on/using hospice services (9937). Optional exclusions include members without a diagnosis of diabetes	Select Health

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
			and diagnosed with polycystic ovarian syndrome (E28.2), gestational (O24.419) or steroid (E09.9) induced diabetes	
Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC)	18-64 years of age	Members between 18 and 64 years of age, with a diagnosis of cardiovascular disease and schizophrenia or schizoaffective disorder, that completed LDL-C testing during the MY	<b>ICD 10 Codes:</b> I20.0, I24.9, I21.4, F20.9, F25.9 <b>CPT Codes:</b> 80061, 83700, 83721 <b>Exclusions:</b> Members on/using hospice services (9937)	Select Health
Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA)	18 years of age and older	Members 18 years of age and older diagnosed with schizophrenia or schizoaffective disorder, that were prescribed an antipsychotic medication, and remained on treatment for at least 80% of the period	<b>ICD 10 Codes:</b> F20.9, F25.9, F25.8 <b>CPT Codes:</b> NA <b>HCPCS Codes:</b> J2794, J0401, J1944, J2798 <b>Exclusions:</b> Members on/using hospice services (9937), members 66 years of age or older residing in a facility, members 66 to 80 years of age with a frailty (L89.000) <i>and</i> advanced illness (A81.00) diagnosis, members 81 and older with a frailty (L89.000) diagnosis, members with a diagnosis of Dementia (F0390) or do not have at least one antipsychotic prescription refill on a different date than initial fill	Select Health, MAP Total, EasyCare and EasyCare Plus
Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM)	1-17 years of age	Members between 1 and 17 years of age, who were prescribed two or more antipsychotic medications and had metabolic testing. <i>Metabolic testing includes blood glucose testing, cholesterol testing or both</i>	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 82465, 80061, 83036, 80047 <b>Exclusions:</b> Members on/using hospice services (9937)	Select Health

**Effectiveness of Care- Care Coordination Measures:**

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Advance Care Planning (ACP)	66 years of age and older	Members between 66 and 80 years of age who have an advanced illness or frailty diagnosis, or receiving palliative care, and members 81 years of age and older who have advanced care planning in the MY	<b>ICD 10 Codes:</b> Z66, G20, G30.9, R41.81 <b>CPT Codes:</b> 99483, 99497 <b>Exclusions:</b> Members on/using hospice services (9937)	MAP Total, EasyCare and EasyCare Plus
Transitions of Care (TRC)	18 years of age and older	Members 18 years of age and older, who have been discharged after an in-patient hospitalization. <i>Rates reported include notification of admission, discharge information, patient engagement post discharge and medication reconciliation post discharge</i>	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 99483, 98966, 99495, 99201 <b>Exclusions:</b> Members on/using hospice services (9937)	MAP Total, EasyCare and EasyCare Plus
Follow-Up After Emergency Department Visit for People with Multiple High-Risk Chronic Conditions (FMC)	18 years of age and older	Members 18 years of age and older, diagnosed with multiple high-risk chronic conditions, who have been in the emergency department (ED) and completed a follow-up appointment within 7 days of the ED visit. <i>High-risk conditions include, but not limited to, COPD, Alzheimer’s disease, chronic kidney disease, heart failure, acute myocardial infarction, atrial fibrillation, and stroke</i>	<b>ICD 10 Codes:</b> J44.9, G30.9, N18.9, I50.9, I21.9, I48.91, I63.9 <b>CPT Codes:</b> 97091, 99366, 99281, 98969, 99201, 98966, 99495 <b>Exclusions:</b> Members on/using hospice services (9937) and members with an ED visit that resulted in an in-patient admission	MAP Total, EasyCare and EasyCare Plus

## HIV Care

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Viral Load Suppression	2 years of age and older	Members 2 years of age and older, who are diagnosed with HIV (Human Immunodeficiency Virus), with their last HIV Viral Load Testing of the year showing a viral load less than 200 copies/mL	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> NA <b>Exclusions:</b> NA	Select Health
Sexually Transmitted Diseases: Chlamydia Screening	13 years of age and older	Members 13 years of age and older, who are diagnosed with HIV (Human Immunodeficiency Virus) who have completed a chlamydia screening within the MY	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> NA <b>Exclusions:</b> Members under 18 years of age, who are not sexually active	Select Health
Sexually Transmitted Diseases: Gonorrhea Screening	13 years of age and older	Members 13 years of age and older, who are diagnosed with HIV (Human Immunodeficiency Virus) who have completed a gonorrhea screening within the MY	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> NA <b>Exclusions:</b> Members under 18 years of age, who are not sexually active	Select Health
Sexually Transmitted Diseases: Syphilis Screening	18 years of age and older, as well as <18 who are sexually active	Members who are sexually active and/or over the age of 18, who are diagnosed with HIV (Human Immunodeficiency Virus) who have completed a serologic syphilis screening within the MY	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> NA <b>Exclusions:</b> Members under 18 years of age, who are not sexually active	Select Health

### Access/Availability of Care Measures:

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Adults' Access to Preventative/Ambulatory Health Services (AAP)	21 years of age and older	Members 20 years of age and older, who have had a preventative care visit within the MY. Preventative visits include ambulatory, online and telephone visits. <i>Age stratifications are 20-44, 45-64 and 65 years of age and older</i>	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 99201, 92002, 98966, 98969 <b>Exclusions:</b> Members on/using hospice services (9937)	Select Health, MAP Total, EasyCare and EasyCare Plus
Annual Dental Visit (ADV)	2-20 years of age	Members between 2 and 20 years of age, who have had at least one dental visit within the MY. <b>Any type of visit with a dentist during the MY meets the criteria.</b> <i>Age stratifications are 2-3, 4-6, 7-10, 11-14, 15-18 and 19-20 years of age</i>	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> NA <b>Exclusions:</b> Members on/using hospice services (9937)	Select Health
Initiation and Engagement of Substance Use Disorder Treatment (IET)	13 years of age and older	Members 13 years of age and older, who have a new alcohol, opioid or other substance abuse disorder, that initiated and obtained treatment. Initiation of SUD treatment within 14 days and engagement of SUD treatment within 34 days of initiation	<b>ICD 10 Codes:</b> HZ2ZZZZ <b>CPT Codes:</b> 99408, 98960, 99212 <b>HCPCS Codes:</b> G2086, G2067, H0020, J2315, H0008 <b>Exclusions:</b> Members on/using hospice services (9937) and any members where the initiation took place in an inpatient admission where the member was discharged after November 27 <sup>th</sup> of MY	Select Health, MAP Total, EasyCare and EasyCare Plus
Prenatal and Postpartum Care (PPC)	None specified	Female members who have had live births on October 8 <sup>th</sup> the year prior through October 7 <sup>th</sup> of the MY. Measure includes timeliness of prenatal care, as well as postpartum care between 7 and 84 days post-delivery <i>Female members who have multiple live births during one pregnancy count once, however</i>	<b>ICD 10 Codes:</b> 10D00Z0 <b>CPT Codes:</b> 88141, 59400, 57170, 59400, 99201, 99500, 98966 <b>HCPCS Codes:</b> G0071, G0463, H1000 <b>Exclusions:</b> Members on/using hospice services (9937), any non-live births and	Select Health

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
		<i>multiple pregnancies resulting in two live deliveries are counted twice</i>	any services provided in an acute inpatient setting (99221)	
Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics <b>(APP)</b>	1-17 years of age	Members between 1 and 17 years of age who were prescribed a new antipsychotic medication, with a documented first-line psycho-social care	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 98960, 90870, 99217, 98969, 90832, 98966, 90791 <b>HCPCS Codes:</b> G0410, G2010, H0002 <b>Exclusions:</b> Members on/using hospice services (9937), members with antipsychotic medication as first-line care	Select Health



### Utilization and Risk Adjusted Utilization Measures:

Measure Name (Measure Code)	Age Criteria	Measure Description	ICD10, CPT Codes and Exclusions	Line of Business (LOB)
Well-Child Visits in the First 30 Months of Life (W30)	31 days to 30 months of age	Members between 31 days old to 15 months of age, who have had 6 or more well-child PCP visits within 15 months, as well as 2 or more well-child PCP visits between 15-30 months.	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 99381-99385, 99391-99395 <b>Exclusions:</b> Members on/using hospice services (9937)	Select Health
Child and Adolescent Well-Care Visits (WCV)	3-21 years of age	Members between 3 and 21 years of age, who have had at least one well-visit PCP or OB/GYN during the MY	<b>ICD 10 Codes:</b> NA <b>CPT Codes:</b> 99381-99385, 99391-99395 <b>Exclusions:</b> Members on/using hospice services (9937)	Select Health

## References and Comments:

- ❖ HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA). Refer to [www.ncqa.org/hedis/](http://www.ncqa.org/hedis/)
- ❖ CPT® (Current Procedural Terminology) is a registered trademark of the American Medical Association
- ❖ HCPCS: Healthcare Common Coding Procedure Coding System Level II
- ❖ LOINC (Logical Observation Identifiers Names and Codes)
- ❖ ICD-10: Centers for Medicare & Medicaid Services at [www.cms.gov/Medicare/Coding/icd10/](http://www.cms.gov/Medicare/Coding/icd10/)
- ❖ For additional information on Advanced Illness and Frailty, refer to <https://www.ncqa.org/blog/improving-care-advanced-illness-frailty/>
- ❖ Quality Assurance Reporting Requirements (QARR) are adopted from NCQA HEDIS with New York state-specific measures added to address health issues of particular importance in New York. This document is not inclusive of all codes and measures. The measure descriptions and codes in this document are derived from the 2022 HEDIS Technical Specifications/Value Set Directory, and the Quality Assurance Reporting Requirements Technical Specifications manual. For the full list of codes, please refer to the NCQA store to purchase.



### **Have questions?**

Please contact us at [QualityManagement@vnsny.org](mailto:QualityManagement@vnsny.org) or [Provider\\_Education@vnsny.org](mailto:Provider_Education@vnsny.org)

For additional information and resources, please visit our provider website and toolkit at <https://www.vnsnychoice.org/for-health-professionals-overview/>